

Hopf-Frobenius algebras and a new Drinfeld double

Joseph Collins¹

Ross Duncan^{1,2}

joseph.collins@strath.ac.uk ross.duncan@strath.ac.uk

¹Department of Computer and Information Sciences
University of Strathclyde
26 Richmond Street, Glasgow, United Kingdom

²Cambridge Quantum Computing Ltd
9a Bridge Street, Cambridge, United Kingdom

The ZX-calculus and related theories are based on so-called interacting Frobenius algebras, where a pair of \dagger -special commutative Frobenius algebras jointly form a pair of Hopf algebras. In this setting we introduce a generalisation of this structure, *Hopf-Frobenius algebras*, starting from a single Hopf algebra which is not necessarily commutative or cocommutative. We provide the necessary and sufficient condition for a Hopf algebra to be a Hopf-Frobenius algebra, and show that every Hopf algebra in $\mathbf{FVect}_{\mathbf{k}}$ is a Hopf-Frobenius algebra. Hopf-Frobenius algebras provide a notion of duality, and give us a “dual” Hopf algebra that is isomorphic to the usual dual Hopf algebra in a compact closed category. We use this isomorphism to construct a Hopf algebra isomorphic to the Drinfeld double that is defined on $H \otimes H$ rather than $H \otimes H^*$.

1 Introduction

In the monoidal categories approach to quantum theory [1, 11] Hopf algebras [29] have a central role in the formulation of complementary observables [10]. In this setting, a quantum observable is represented as special commutative \dagger -Frobenius algebra; a pair of such observables are called *strongly complementary* if the algebra part of the first and the coalgebra part of the second jointly form a Hopf algebra. In abstract form, this combination of structures has been studied under the name “interacting Frobenius algebras” [14] where it is shown that relatively weak commutation rules between the two Frobenius algebras produce the Hopf algebra structure. From a different starting point Bonchi et al [5] showed that a distributive law between two Hopf algebras yields a pair of Frobenius structures, an approach which has been generalised to provide a model of Petri nets [4]. Given the similarity of the two structures it is appropriate to consider both as exemplars of a common family of *Hopf-Frobenius algebras*.

In the above settings, the algebras considered are both commutative and cocommutative. However more general Hopf algebras, perhaps not even symmetric, are a ubiquitous structure in mathematical physics, finding application in gauge theory [24], condensed matter theory [27], quantum field theory [6] and quantum gravity [23]. In this paper we take the first steps towards generalising the concept of Hopf-Frobenius algebra to the non-commutative case, and opening the door to applications of categorical quantum theory in other areas of physics.

Loosely speaking, a Hopf-Frobenius algebra consists of two monoids and two comonoids such that one way of pairing a monoid with a comonoid gives two Frobenius algebras, and the other pairing yields two Hopf algebras, with the additional condition that antipodes are constructed from the Frobenius forms. This schema is illustrated in Figure 1. In Section 3 we give the precise definition of Hopf-Frobenius algebras and state the necessary and sufficient conditions to extend




Figure 1: The elements of a Hopf-Frobenius algebra

a Hopf algebra to a Hopf-Frobenius algebra in an arbitrary symmetric monoidal category. It was previously known that in \mathbf{FVect}_k , the category of finite dimensional vector spaces, every Hopf algebra carries a Frobenius algebra on both its monoid [22] and its comonoid [12, 20]; in fact every Hopf algebra in \mathbf{FVect}_k is Hopf-Frobenius. In Section 4 we briefly present some examples which are not abelian group algebras. In Section 5 we show the structure of a Hopf-Frobenius can be used to give a simpler version of the Drinfeld double construction.

2 Preliminaries

We assume that the reader is familiar with strict symmetric monoidal categories and their diagrammatic notation; see Selinger [26] for a thorough treatment. We make the convention that diagrams are read from top to bottom. Since we work in categories with duals, the wires carry an orientation: downwards for the original object, upwards for its dual. We will otherwise omit the object names from the wires except where doing so would create ambiguity.

Definition 2.1. In a monoidal category \mathcal{C} with objects A and B , B is *left dual* to A if there exist morphisms $d : I \rightarrow A \otimes B$ and $e : B \otimes A \rightarrow I$ such that

$$\begin{array}{c} \text{cap} \\ \downarrow \\ A \end{array} = \begin{array}{c} \text{cup} \\ \downarrow \\ A \end{array} \quad \text{and} \quad \begin{array}{c} \text{cup} \\ \downarrow \\ B \end{array} = \begin{array}{c} \text{cap} \\ \downarrow \\ B \end{array}$$


In this circumstance A is *right dual* to B . Note that if \mathcal{C} is symmetric then left duals and right duals coincide.

The morphisms d and e are usually called the unit and counit; for reasons which will become obvious shortly we avoid that terminology and refer to them as the *cap* and the *cup*. Note that if an object has a dual it is unique up to isomorphism (see Lemma C.1).

Definition 2.2. A *compact closed category* [19] is symmetric monoidal category where every object A has a chosen right dual (A^*, d_A, e_A) . In the graphical notation we depict the cup and cap in the obvious way:


$$d_A := \begin{array}{c} \text{cap} \\ \downarrow \\ A^* \end{array} \quad e_A := \begin{array}{c} \text{cup} \\ \downarrow \\ A \end{array}$$

Proposition 2.3 ([19]). *Let \mathcal{C} be a compact closed category. By defining $f^* : B^* \rightarrow A^*$ as*



the assignment of duals $A \mapsto A^$ extends uniquely to a strong monoidal functor $(\cdot)^* : \mathcal{C}^{\text{op}} \rightarrow \mathcal{C}$, with natural isomorphisms $(A \otimes B)^* \cong B^* \otimes A^*$, $A^{**} \cong A$, and $I^* \cong I$ and, further, d and e are natural transformations.*


Since we have the freedom to choose, we will take the isomorphisms listed above as the definition of the dual. This means that the cap for tensor products of objects is “nested”, as shown below,



and similarly for the cup.

The main foci of this work – Frobenius and Hopf algebras – combine the structure of a monoid and a comonoid on the same object. See Appendix C.2 for basic definitions.

Definition 2.4. A *Frobenius algebra* in a symmetric monoidal category \mathcal{C} consists of a monoid and a comonoid on the same object, obeying the Frobenius law, shown below on the left:



A Frobenius algebra is called *special* or *separable* when it obeys the equation above right, and *quasi-special* when it obeys the special equation up to a scalar factor. A Frobenius algebra is commutative when its monoid is, and cocommutative when its comonoid is.

Lemma 2.5. *Every Frobenius algebra induces a cup and a cap which make the object self-dual.*


Proof. Given the Frobenius algebra $(\curvearrowright, \circlearrowleft, \curvearrowright, \circlearrowright)$ define the cup and cap as shown below.



From here the snake equation follows directly. □

Definition 2.4, due to Carboni and Walters [7], has a pleasing symmetry between the monoid and comonoid parts. However, an older equivalent definition will be useful in later sections¹.

Definition 2.6. A *Frobenius algebra* in a symmetric monoidal category \mathcal{C} consists of a monoid $(F, \curvearrowright, \circlearrowleft)$ and a *Frobenius form* $\beta : F \otimes F \rightarrow I$, which admits an inverse, $\bar{\beta} : I \rightarrow F \otimes F$, satisfying:




¹See Fauser’s survey [15] for several equivalent definitions.

To see that Definition 2.4 implies this definition it suffices to take the cup and cap defined above as β and $\bar{\beta}$. The converse implication is left as an exercise.

Frobenius forms are far from unique: there is one for each invertible element of the monoid (see AppendixC.3).

Special Frobenius algebras can be understood as arising from a distribution law of comonoids over monoids [21]. In the other direction, distributing monoids over comonoids yields bialgebras. *NOTE.* Unlike the preceding section, in our discussion of bialgebras and Hopf algebras, we will use different colours for the monoid and comonoid parts of the structure.


Definition 2.7. A *bialgebra* in symmetric monoidal category \mathcal{C} consists of a monoid and a comonoid on the same object, which jointly obey the *copy*, *cocopy*, *bialgebra*, and *scalar* laws depicted below.



Note that the dashed box above represents an empty diagram. A *bialgebra morphism* is an endomorphism of the object which commutes with both the monoid and the comonoid structures.

Remark 2.8. Some works, notably on the ZX-calculus [10, 2, 17] and related theories [14], the last axiom is dropped and the other equations modified by a scalar factor, to give a *scaled bialgebra*. Here we use the standard definition: The Frobenius algebras we construct will not be special.

Definition 2.9. A *Hopf algebra* consists of a bialgebra $(H, \overset{\circ}{\circlearrowleft}, \overset{\circ}{\circlearrowright}, \bullet, \bullet)$ and an endomorphism $s : H \rightarrow H$ called the *antipode* which satisfies the *Hopf law*:




Where unambiguous, we abuse notation slightly and use H to refer the whole Hopf algebra. Following Street [28], we can define another Hopf algebra H^{op} on the same object, having the same unit and counit, but with the arguments of the multiplication and comultiplication swapped:



Replacing only the comultiplication as above yields a bialgebra H^σ which is not necessarily Hopf. We quote the following basic properties from Street [28].


Proposition 2.10. For a Hopf algebra H :

1. The antipode s is unique.
2. $s : H^{\text{op}} \rightarrow H$ is a bialgebra homomorphism, i.e.



3. H^σ is a Hopf algebra if and only if s is invertible, in which case the antipode of H^σ is s^{-1} .
4. If H is commutative or cocommutative then $s \circ s = \text{id}_H$.

Definition 2.11. Let $(H, \begin{smallmatrix} \curvearrowright \\ \circlearrowleft \end{smallmatrix}, \begin{smallmatrix} \circlearrowright \\ \curvearrowleft \end{smallmatrix}, \begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}, \blacksquare)$ be a Hopf algebra, and suppose that the object H has a left dual H^* . We define the *dual Hopf algebra* $(H^*, \begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}^*, \begin{smallmatrix} \curvearrowright \\ \circlearrowleft \end{smallmatrix}^*, \begin{smallmatrix} \circlearrowright \\ \curvearrowleft \end{smallmatrix}^*, \blacksquare^*)$ as :



It's straightforward to prove that H^* is indeed a Hopf algebra using the equations of Def 2.1. In later sections it will be helpful to consider duals with respect to different cups and caps, in which case we will vary notation accordingly but the same construction is used in all cases.

3 Hopf-Frobenius Algebras

We now arrive at the main subject of this paper, Hopf-Frobenius algebras in an arbitrary symmetric monoidal category \mathcal{C} . These algebras generalise interacting Frobenius algebras [10, 14], and share the same gross structure. It will be helpful to introduce a weaker notion first.

Definition 3.1. A *pre-Hopf-Frobenius algebra* or *pre-HF algebra* consists of an object H bearing a green monoid $(\begin{smallmatrix} \curvearrowright \\ \circlearrowleft \end{smallmatrix}, \begin{smallmatrix} \circlearrowright \\ \curvearrowleft \end{smallmatrix})$, a green comonoid $(\begin{smallmatrix} \circlearrowright \\ \curvearrowleft \end{smallmatrix}, \begin{smallmatrix} \circlearrowleft \\ \curvearrowright \end{smallmatrix})$, a red monoid $(\begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}, \begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix})$, a red comonoid $(\begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}, \begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix})$ and an endomorphism such \blacksquare that $(\begin{smallmatrix} \curvearrowright \\ \circlearrowleft \end{smallmatrix}, \begin{smallmatrix} \circlearrowright \\ \curvearrowleft \end{smallmatrix}, \begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}, \begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix})$ and $(\begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}, \begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}, \begin{smallmatrix} \curvearrowright \\ \circlearrowleft \end{smallmatrix}, \begin{smallmatrix} \circlearrowright \\ \curvearrowleft \end{smallmatrix})$ are Frobenius algebras, and $(\begin{smallmatrix} \curvearrowright \\ \circlearrowleft \end{smallmatrix}, \begin{smallmatrix} \circlearrowright \\ \curvearrowleft \end{smallmatrix}, \begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}, \begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}, \blacksquare)$ is a Hopf algebra.

Definition 3.2. A *Hopf-Frobenius algebra*, or *HF algebra*, is a pre-Hopf-Frobenius algebra where \blacksquare satisfies the left equation below,



and with \blacksquare defined as in the right equation above, $(\begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}, \begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}, \begin{smallmatrix} \curvearrowright \\ \circlearrowleft \end{smallmatrix}, \begin{smallmatrix} \circlearrowright \\ \curvearrowleft \end{smallmatrix}, \blacksquare)$ is a Hopf algebra.

We refer to the four algebras that make up an HF algebra by the colour² of their *multiplication*, so that $(\begin{smallmatrix} \curvearrowright \\ \circlearrowleft \end{smallmatrix}, \begin{smallmatrix} \circlearrowright \\ \curvearrowleft \end{smallmatrix}, \begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}, \begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}, \blacksquare)$ is the *green* Hopf algebra, $(\begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}, \begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}, \begin{smallmatrix} \curvearrowright \\ \circlearrowleft \end{smallmatrix}, \begin{smallmatrix} \circlearrowright \\ \curvearrowleft \end{smallmatrix}, \blacksquare)$ is the *red* Frobenius algebra, etc.

Remark 3.3. Despite their distinct definitions, the two antipodes may coincide; for example if both Frobenius algebras are symmetric, as in a group algebra, then they are equal.

We now move on to the main topic of the section: under what conditions does a Hopf algebra extend to a Hopf-Frobenius algebra? Henceforward, unless otherwise stated, \mathcal{C} will denote a symmetric monoidal category, and H will denote a Hopf algebra $(H, \begin{smallmatrix} \curvearrowright \\ \circlearrowleft \end{smallmatrix}, \begin{smallmatrix} \circlearrowright \\ \curvearrowleft \end{smallmatrix}, \begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}, \begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}, \blacksquare)$ in \mathcal{C} . Omitted proofs are found in Appendix A.

A key concept is that of an integral. Pareigis [25] proved³ that in $\mathbf{FPMo d}_{\mathbf{R}}$, the category of finitely generated projective modules over a commutative ring, a Hopf algebra has Frobenius

²If you are reading this document in monochrome *green* will appear as light grey and *red* as dark grey.

³This is a generalisation of earlier work by Larson and Sweedler [22] showing that the space of integrals in \mathbf{FVect}_k is always isomorphic to k .

structure when its space of integrals is isomorphic to the ring. We will proceed in a similar direction.

Definition 3.4. A *left (co)integral* on H is a copoint $\downarrow : H \rightarrow I$ (resp. a point $\uparrow : I \rightarrow H$), satisfying the equations:

A *right (co)integral* is defined similarly.

Definition 3.5. An *integral Hopf algebra* $(H, \uparrow, \downarrow)$ is a Hopf algebra H equipped with a choice of right integral \downarrow , and left cointegral \uparrow , such that $\downarrow \circ \uparrow = \text{id}_I$.

Lemma 3.6. Let $(H, \uparrow, \downarrow)$ be an integral Hopf algebra. Then the following map is the inverse of the antipode.

Lemma 3.7. Let $(H, \uparrow, \downarrow)$ be an integral Hopf algebra, and define

then β is a Frobenius form for $(H, \uparrow, \downarrow)$ iff γ is a right inverse for β .

Per Definition 2.6, \downarrow is the counit of this Frobenius algebra and the comultiplication is obtained by dualising \uparrow with β .

Lemma 3.7 shows an example of a duality structure which produces a Frobenius algebra from an integral Hopf algebra. We wish to treat all such dualities in a uniform way, and also to weaken the requirements on such a duality. For this purpose, we shall define the following concept.

Definition 3.8. Let A and B be objects in a symmetric monoidal category \mathcal{C} . A is a *right half dual* of B if there exist morphisms $\frown : I \rightarrow A \otimes B$ and $\smile : B \otimes A \rightarrow I$ which satisfy the left equation of 2.1. In this circumstance, B is a *left half dual* of A .


Half duals are a strict generalisation of duals in the sense of Definition 2.1. Further, any integral Hopf algebra $(H, \uparrow, \downarrow)$ makes H left half dual to itself, via the morphisms β and γ of Lemma 3.7. Unlike true duals, an object may have non-isomorphic half duals. For example, if B is left dual to A , with a retract $m : B \hookrightarrow C$ for some section $m' : C \rightarrow B$, then C is also left half dual of A .

Definition 3.9. Let the object H have a right half dual H^* . The *integral section* $\mathcal{I} : H \rightarrow H$ is defined as shown below.

Note that this definition does *not* depend on the choice of half dual – see Lemma A.1

In $\mathbf{FPMod}_{\mathbf{R}}$, \mathcal{I} is the section of the natural injection from the space of left integrals into H . As such, it acts trivially on integrals, and for every element $v \in H$, $\mathcal{I}(v)$ is either a left integral or 0. In Lemma A.3 we show that this holds in the general case.

Definition 3.10. We say that a Hopf algebra satisfies the *Frobenius condition* if there exists maps \uparrow and \downarrow such that




Theorem 3.11. If H satisfies the Frobenius condition, then H is a pre-HF algebra with the Frobenius forms and their inverses as shown below.



Further, $(H, \uparrow, \downarrow)$ is an integral Hopf algebra.

The explicit definitions of the green comonoid and red monoid structures are shown below.



Corollary 3.12. Every Hopf algebra which satisfies the Frobenius condition defines a pre-HF algebra.

We can rephrase Pareigis' condition as the existence of an integral \uparrow in H such that each other integral is equal to a scalar multiple of \uparrow .

In a monoidal category, an object A is said to have *enough points* if, for all morphisms $f, g : A \rightarrow B$, we have

$$(\forall x : I \rightarrow A, \quad fx = gx) \Rightarrow f = g .$$

Lemma 3.13. If H satisfies the Frobenius condition then every left cointegral (right integral) is a scalar multiple of \uparrow (resp. \downarrow); further if H has enough points then the converse holds.

Since $\mathbf{FPMod}_{\mathbf{R}}$ (and $\mathbf{FVect}_{\mathbf{k}}$) are categories where every object has enough points, Lemma 3.13 implies Pareigis' condition is exactly the Frobenius condition.

We now consider the question of when a pre-HF algebra is Hopf-Frobenius. There are two issues to address: whether the red Hopf algebra exists, and whether the two antipodes have the required form.

Lemma 3.14. Let H be a pre-HF algebra; $(H, \uparrow, \downarrow)$ is an integral Hopf algebra if and only if $\square = \text{diagram}$.

Lemma 3.15. Let H be a pre-HF algebra such that $(H, \uparrow, \downarrow)$ is an integral Hopf algebra, and let $(\cdot)^\circ$ be the duality defined by the green Frobenius algebra (cf. Lemma 2.5). Then:



Corollary 3.16. *Let H be a pre-HF algebra such that (H, \bullet, \circ) is an integral Hopf algebra. $(H, \overleftarrow{\bullet}, \bullet, \overleftarrow{\circ}, \circ, \blacksquare)$ forms a Hopf algebra, where $\blacksquare = \overleftarrow{\circ} \bullet$.*

Putting all of the above together, we obtain the main result of this section.

Theorem 3.17. *Let H be a Hopf algebra such that the object H has some weak right dual H^* . Then H is a Hopf-Frobenius algebra if and only if H fulfills the Frobenius condition.*

Proof. By Theorem 3.11, the Frobenius condition implies the existence of the two Frobenius algebras, and that H is an integral Hopf algebra. By Lemma 3.13 we have that (H, \bullet, \circ) is integral, hence by Lemma 3.14 that \blacksquare has the required form. Corollary 3.16 then completes the proof. For the converse direction, it is straightforward to show that every Hopf-Frobenius algebra satisfies the Frobenius condition. \square

One might wonder whether there is any gap between a pre-HF algebra and an HF algebra. The requirement on the form of the antipodes is decisive here. Suppose that we have an HF algebra H . Formally exchanging the colours yields a pre-HF algebra, with two Hopf algebras. However, unless the Frobenius algebras are symmetric, this transformation does not preserve the form of the antipodes hence the resulting structure is not an HF algebra. However the gap is a narrow one, as the following result shows.

Theorem 3.18. *Let H be a pre-HF algebra such that, for some \blacksquare , $(H, \overleftarrow{\bullet}, \bullet, \overleftarrow{\circ}, \circ, \blacksquare)$ forms a Hopf algebra; then H extends to a Hopf-Frobenius algebra.*

4 Examples

Combined with the results of Larson and Sweedler [22], Pareigis [25], and Lemma 3.13, Theorem 3.17 implies that any Hopf algebra in \mathbf{FVect}_k is Hopf-Frobenius. This allows the direct extension of [14] to non-abelian group algebras, but there are plenty of other examples. We briefly mention some examples which are neither commutative nor cocommutative.

Example 4.1. Let k be a field with a primitive n^{th} root of unity z . The Taft Hopf algebras [30] are a family of Hopf algebras in \mathbf{FVect}_k whose antipodes have order $2n$. Generically, the algebra $(H, \mu, 1, \Delta, \epsilon, s)$ is generated by elements x and g , such that $x^n = 0$, $g^n = 1$, and $gx = zxg$. The coalgebra is defined $\Delta(x) = 1 \otimes x + x \otimes g$, and $\Delta(g) = g \otimes g$, with $\epsilon(x) = 0$ and $\epsilon(g) = 1$. The antipode is $s(x) = -xg^{-1}$, $s(g) = g$, and the rest of the structure follows from the Hopf algebra axioms. We may see that H has the basis $x^\alpha g^\beta$, where $0 \leq \alpha, \beta \leq n-1$, so this will imply that H is n^2 dimensional. One can calculate that the left integral of H is

$$\sum_{i=1}^n z^{-i} g^i x^{n-1}$$

and the right cointegral is the functional that takes x^{n-1} to 1 and every other basis element to 0. We explicitly construct the HF algebra of the Taft Hopf algebra when $n = 2$ in the appendix.

Example 4.2. Hopf algebras which arise as the quantum enveloping algebra of Lie algebras are type of quantum group. Since these are infinite dimensional, they cannot be Hopf-Frobenius algebras. However their finite dimensional quotients will be Hopf-Frobenius. See Kassel [18] for an example.

Moving away from \mathbf{FVect}_k , we consider \mathbf{Rel} , the category of sets and relations.

Example 4.3. Let G be an infinite group. Following Hasegawa [16] we can construct its group algebra in \mathbf{Rel} . The integral is $\{(\star, g) \mid g \in G\}$ and the cointegral is the singleton $(1, \star)$. The construction detailed in Theorem 3.11 recovers the expected multiplication and comultiplication relations:

$$\begin{aligned} \text{green circle} &:= a \mapsto (b, c) \text{ such that } a = bc \\ \text{red circle} &:= (a, b) \mapsto \begin{cases} a & \text{if } a = b \\ \emptyset & \text{otherwise} \end{cases} \end{aligned}$$


We look forward to discovering more exotic examples.

5 A new Drinfeld double

Braided categories of modules over a Hopf algebra are widely used in physics, where they give solutions to the Yang-Baxter equation and in low dimensional topology, where they are used to find invariants. However the category of modules over a Hopf algebra is braided if and only if the Hopf algebra is *quasi-triangular*. The *Drinfeld double* [13] is a construction that takes a Hopf algebra H in \mathbf{FVect}_k , and produces a quasi-triangular Hopf algebra $D(H)$ on the object $H \otimes H^*$. In this section we use the self-duality of a Hopf-Frobenius algebra to construct the canonical isomorphism $H \cong H^*$ and thus define a simpler version of the Drinfeld double on $H \otimes H$.

We will assume that \mathcal{C} is a compact closed category. We denote the green and red Hopf algebras of H as H_\circ and H_\bullet respectively. We use the generalisation of Drinfeld’s original construction to symmetric monoidal categories, due to Chen [8].


Definition 5.1. Let H be a HF algebra on \mathcal{C} . By Proposition C.1, we may define an isomorphism $\square: H \rightarrow H^*$, with inverse $\square^{-1}: H^* \rightarrow H$ as



Lemma 5.2. The morphism \square is a Hopf algebra homomorphism between H_\bullet^{op} and H_\circ^* .

Definition 5.3. A Hopf algebra H is *quasi-triangular* there exists a *universal R-matrix* $R: I \rightarrow H \otimes H$ such that


- R is invertible with respect to green circle




All cocommutative Hopf algebras are quasi-triangular, with $\eta \otimes \eta$ as the universal R -matrix. This definition is motivated by the following theorem [18].

Theorem 5.4. *The category of modules over a Hopf algebra is braided if and only if the Hopf algebra is quasi-triangular*


Definition 5.5. Let H be a Hopf algebra in \mathcal{C} with an invertible antipode. The *Drinfeld double* of H , denoted $D(H) = (H \otimes H^*, \mu, 1, \Delta, \epsilon, s)$, is a Hopf algebra defined in the following manner:




Theorem 5.6 (Drinfeld[13, 8]). *$D(H)$ is quasi-triangular, with the universal R-matrix shown below.*




Now, consider the composite of the map $1 \otimes \square$ with the multiplication of the Drinfeld double:




Similarly, composing the universal R-matrix with the inverse isomorphism, $1 \otimes \square^{-1}$ gives us



Definition 5.7. Let H be a HF algebra. The *red Drinfeld double*, denoted $D_{\bullet}(H) = (H \otimes H^*, \mu_{\bullet}, 1_{\bullet}, \Delta_{\bullet}, \epsilon_{\bullet}, s_{\bullet})$, is a Hopf algebra on the object $H \otimes H$ with structure maps



Corollary 5.8. $D_\bullet(H)$ is a quasi-triangular Hopf algebra isomorphic to the Drinfeld double, with universal R -matrix



6 Conclusion and further work

We have generalised the notions of interacting Frobenius algebras [10, 14] and interacting Hopf algebras [5] to the non-commutative case, and in the process shown that they are rather common structures. This work could be viewed as an extension of classical results showing that concrete Hopf algebras over finite dimensional vector spaces are also Frobenius algebras [22]. Another perspective is that we make precise how much ambient symmetry is required to obtain a Hopf-Frobenius algebra. The original setting of interacting Frobenius algebras [10] was a \dagger -compact category, which provides a lot of duality on top of the commutative algebras themselves. We show that none of this structure is necessary: all that is required is one-sided *half-dual* for the carrier object. The major question that remains is to pin down exactly when the Frobenius condition holds; as Lemma 3.13 shows, this is tightly related to the existence of integrals. Compact closure does not suffice to guarantee this: in $\mathbf{FPMod}_{\mathbf{R}}$ there are Hopf algebras which are not Frobenius.

While we have established that Hopf algebras are frequently Hopf-Frobenius, the resulting Frobenius algebras need not be well behaved (commutative, dagger, special) as in the original quantum setting [9]. It remains to investigate what Frobenius structures arise from “interesting” Hopf algebras, and whether they have any application in the categorical quantum mechanics programme, or conversely, how HF algebras may be applied in the study of quantum groups. Weaker structures such as the ill-named coFrobenius algebras or the stateful resource calculus of Bonchi et al [4] perhaps offer a new route to study infinite dimensional systems. Beyond this, natural generalisations to the braided or planar cases suggest themselves, although this will push diagrammatic reasoning to its limits.

Our new Drinfeld double construction suggests that HF algebras could find applications in topological quantum computation, particularly for error correcting codes, an area where the ZX-calculus is already used [3]. The smallest non-abelian group is S_3 , whose group algebra fits in 3 qubits with room to spare.

References

- [1] S. Abramsky & B. Coecke (2004): *A categorical semantics of quantum protocols*. In: *Proceedings of the 19th Annual IEEE Symposium on Logic in Computer Science: LICS 2004*, IEEE Computer Society, pp. 415–425.
- [2] Miriam Backens (2015): *Making the stabilizer ZX-calculus complete for scalars*. In Chris Heunen, Peter Selinger & Jamie Vicary, editors: *Proceedings of the 12th International Workshop on Quantum Physics and Logic (QPL 2015)*, *Electronic Proceedings in Theoretical Computer Science* 195, pp. 17–32, doi:10.4204/EPTCS.195.2.
- [3] Niel de Beaudrap & Dominic Horsman (2017): *The ZX calculus is a language for surface code lattice surgery*. In: *Proc. QPL2017*.
- [4] Filippo Bonchi, Joshua Holland, Robin Piedeleu, Paweł Sobociński & Fabio Zanasi (2019): *Diagrammatic Algebra: From Linear to Concurrent Systems*. *Proceedings of the ACM on Programming*

- Languages* 3(POPL), pp. 25:1–25:28, doi:10.1145/3290338. Available at <http://doi.acm.org/10.1145/3290338>.
- [5] Filippo Bonchi, Paweł Sobociński & Fabio Zanasi (2017): *Interacting Hopf Algebras*. *Journal of Pure and Applied Algebra* 221(1), pp. 144–184.
- [6] Christian Brouder (2009): *Quantum field theory meets Hopf algebra*. *Mathematische Nachrichten* 282(12), pp. 1664–1690, doi:10.1002/mana.200610828. Available at <https://onlinelibrary.wiley.com/doi/abs/10.1002/mana.200610828>.
- [7] A. Carboni & R.F.C. Walters (1987): *Cartesian bicategories I*. *Journal of Pure and Applied Algebra* 49(1-2), doi:10.1016/0022-4049(87)90121-6.
- [8] Hui-Xiang Chen (2000): *Quantum doubles in monoidal categories*. *Communications in Algebra* 28(5), pp. 2303–2328.
- [9] B. Coecke, D. Pavlovic & J. Vicary (2013): *A new description of orthogonal bases*. *Math. Structures in Comp. Sci.* 23(3), pp. 555–567, doi:10.1017/S0960129512000047.
- [10] Bob Coecke & Ross Duncan (2011): *Interacting Quantum Observables: Categorical Algebra and Diagrammatics*. *New J. Phys* 13(043016), doi:10.1088/1367-2630/13/4/043016. Available at <http://iopscience.iop.org/1367-2630/13/4/043016/>.
- [11] Bob Coecke & Aleks Kissinger (2017): *Picturing Quantum Processes: A First Course in Quantum Theory and Diagrammatic Reasoning*. Cambridge University Press.
- [12] Yukio Doi & Mitsuhiro Takeuchi (2000): *Bi-Frobenius algebras*. In Nicolás Andruskiewitsch, Walter Ricardo Ferrer Santos & Hans-Jürgen Schneider, editors: *New trends in Hopf algebra theory, Contemporary Mathematics* 267, American Mathematical Society, pp. 67–98, doi:10.1090/conm/267/04265.
- [13] Vladimir Gershonovich Drinfeld (1986): *Quantum groups*. *Zapiski Nauchnykh Seminarov POMI* 155, pp. 18–49.
- [14] Ross Duncan & Kevin Dunne (2016): *Interacting Frobenius Algebras are Hopf*. In Martin Grohe, Eric Koskinen & Natarajan Shankar, editors: *Proceedings of the 31st Annual ACM/IEEE Symposium on Logic in Computer Science, LICS '16, New York, NY, USA, July 5-8, 2016*, LICS '16, ACM, pp. 535–544, doi:10.1145/2933575.2934550.
- [15] Bertfried Fauser (2013): *Some Graphical Aspects of Frobenius Algebras*. In Chris Heunen, Mehrnoosh Sadrzadeh & Edward Grefenstette, editors: *Quantum Physics and Linguistics: A Compositional, Diagrammatic Discourse*, Oxford, doi:10.1093/acprof:oso/9780199646296.003.0002.
- [16] Masahito Hasegawa (2010): *Bialgebras in rel*. *Electronic Notes in Theoretical Computer Science* 265, pp. 337–350.
- [17] Emmanuel Jeandel, Simon Perdrix & Renaud Vilmart (2017): *A Complete Axiomatisation of the ZX-Calculus for Clifford+T Quantum Mechanics*. In: *LICS '18- Proceedings of the 33rd Annual ACM/IEEE Symposium on Logic in Computer Science*, arXiv:1705.11151, ACM, doi:10.1145/3209108.3209131.
- [18] Christian Kassel (2012): *Quantum groups*. 155, Springer Science & Business Media.
- [19] G.M. Kelly & M.L. Laplaza (1980): *Coherence for Compact Closed Categories*. *Journal of Pure and Applied Algebra* 19, pp. 193–213.
- [20] M. Koppinen (1996): *On Algebras with Two Multiplications, Including Hopf Algebras and Bose–Mesner Algebras*. *Journal of Algebra* 182(1), pp. 256 – 273, doi:10.1006/jabr.1996.0170. Available at <http://www.sciencedirect.com/science/article/pii/S0021869396901702>.
- [21] Stephen Lack (2004): *Composing PROPs*. *Theory and Applications of Categories* 13(9), pp. 147–163.
- [22] Richard Gustavus Larson & Moss Eisenberg Sweedler (1969): *An Associative Orthogonal Bilinear Form for Hopf Algebras*. *American Journal of Mathematics* 91(1), pp. 75–94. Available at <https://www.jstor.org/stable/2373270>.

- [23] S Majid (1988): *Hopf algebras for physics at the Planck scale*. *Classical and Quantum Gravity* 5(12), pp. 1587–1606, doi:10.1088/0264-9381/5/12/010. Available at <https://doi.org/10.1088/0264-9381/5/12/010>.
- [24] Catherine Meusburger (2017): *Kitaev Lattice Models as a Hopf Algebra Gauge Theory*. *Communications in Mathematical Physics* 353(1), pp. 413–468, doi:10.1007/s00220-017-2860-7. Available at <https://doi.org/10.1007/s00220-017-2860-7>.
- [25] Bodo Pareigis (1971): *When Hopf algebras are Frobenius algebras*. *Journal of Algebra* 18(4), pp. 588 – 596, doi:10.1016/0021-8693(71)90141-4. Available at <http://www.sciencedirect.com/science/article/pii/0021869371901414>.
- [26] Peter Selinger (2011): *A survey of graphical languages for monoidal categories*. In Bob Coecke, editor: *New structures for physics, Lecture Notes in Physics* 813, Springer, pp. 289–355.
- [27] J.K. Slingerland & F.A. Bais (2001): *Quantum groups and non-Abelian braiding in quantum Hall systems*. *Nuclear Physics B* 612(3), pp. 229 – 290, doi:[https://doi.org/10.1016/S0550-3213\(01\)00308-X](https://doi.org/10.1016/S0550-3213(01)00308-X). Available at <http://www.sciencedirect.com/science/article/pii/S055032130100308X>.
- [28] R. Street (2007): *Quantum Groups: A Path to Current Algebra*. Australian Mathematical Society Lecture Series, Cambridge University Press.
- [29] Moss E. Sweedler (1969): *Hopf Algebras*. W. A. Benjamin Inc.
- [30] Earl J. Taft (1971): *The Order of the Antipode of Finite-dimensional Hopf Algebra*. *Proceedings of the National Academy of Sciences* 68(11), pp. 2631–2633, doi:10.1073/pnas.68.11.2631. Available at <https://www.pnas.org/content/68/11/2631>.

A Proofs omitted from the main body of the paper

Lemma 3.6. *Let $(H, \uparrow, \downarrow)$ be an integral Hopf algebra. Then the following map is the inverse of the antipode.*

$$\square^{-1} := \begin{array}{c} \uparrow \\ \bullet \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \bullet \\ \uparrow \end{array}$$

Proof. From the definition of \square^{-1} , we see that

$$\begin{array}{c} \uparrow \\ \bullet \\ \downarrow \end{array} \square^{-1} = \begin{array}{c} \uparrow \\ \bullet \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \bullet \\ \uparrow \end{array} = \begin{array}{c} \downarrow \\ \bullet \\ \uparrow \end{array} = \begin{array}{c} \downarrow \\ \bullet \\ \uparrow \end{array} = \begin{array}{c} \downarrow \\ \bullet \\ \uparrow \end{array}$$

This implies that H^σ is a Hopf algebra. Recall that by Proposition 2.10, when H^σ is a Hopf algebra, then the antipode of H^σ is the inverse of the antipode of H . Thus, $\square^{-1} \circ \square = 1$ \square

Lemma 3.7. *Let $(H, \uparrow, \downarrow)$ be an integral Hopf algebra, and define*

$$\beta := \begin{array}{c} \downarrow \\ \downarrow \\ \bullet \\ \downarrow \end{array} \quad \gamma := \begin{array}{c} \uparrow \\ \bullet \\ \square \\ \downarrow \end{array}$$

then β is a Frobenius form for $(H, \uparrow, \downarrow)$ iff γ is a right inverse for β .

Proof. Note that γ is left inverse to β by Lemma 3.6; hence if it is also right inverse then the conditions of Definition 2.6 are satisfied. Conversely, suppose that β is a Frobenius form; then there exists some $\bar{\beta}$ such that

$$\begin{array}{c} \bar{\beta} \\ \downarrow \\ \bullet \\ \downarrow \end{array} = \downarrow$$


Appealing again to Lemma 3.6 we have

$$\begin{array}{c} \bar{\beta} \\ \downarrow \\ \downarrow \end{array} = \begin{array}{c} \uparrow \\ \bullet \\ \square \\ \downarrow \end{array} \begin{array}{c} \bar{\beta} \\ \downarrow \\ \downarrow \end{array} = \begin{array}{c} \uparrow \\ \bullet \\ \square \\ \downarrow \end{array}$$

hence, γ is the right inverse of β . \square


Lemma A.1. *When H has two half dual structures, \uparrow, \downarrow and \uparrow', \downarrow' , then the integral sections coincide.*

Proof.




□

Lemma A.2.




Proof.



□

other direction, let \uparrow be a left cointegral. Then




The proof for integrals is similar. □

Theorem 3.11. *If H satisfies the Frobenius condition, then H is a pre-HF algebra with the Frobenius forms and their inverses as shown below.*




Further, $(H, \uparrow, \downarrow)$ is an integral Hopf algebra.

Proof. First off, note that the Frobenius condition implies that $\mathcal{I} \circ \uparrow = \uparrow$, and $\downarrow \circ \mathcal{I} = \downarrow$. Hence, by Lemma A.3, \uparrow is a left cointegral and \downarrow is a right integral. Thus, $(H, \uparrow, \downarrow)$ is an integral Hopf algebra. We may now show that the following maps cancel out:




where (1.) is due to the Frobenius condition, (2.) comes from associativity, and (3.) is due to the Hopf law. We then get the following identity



Thus, by Lemma 3.7 we have our result. □

Lemma 3.13. (forward direction) *Let H be a Hopf algebra that fulfills the Frobenius condition. Every left cointegral (right integral) is a scalar multiple of \blacktriangleright (resp. \circlearrowleft).*


Proof. Let \blacktriangleright be a left cointegral on H . Then by Lemma A.1, as $\blacktriangleright = \blacktriangleright \circ \mathcal{I}$




The proof for right integrals is similar. □

Lemma 3.13. (backward direction) *Let H have enough points, and $(H, \blacktriangleright, \blacktriangleleft)$ be an integral Hopf algebra. Suppose that every left cointegral (right integral) is a scalar multiple of \blacktriangleright (resp. \blacktriangleleft). Then H fulfills the Frobenius condition.*


Proof. By Lemma A.3, for all points a (copoint b), $\mathcal{I} \circ a$ is a cointegral and $b \circ \mathcal{I}$ is an integral. Then, by hypothesis



for some scalar k . Since \blacktriangleleft is an integral, $\blacktriangleleft \circ \mathcal{I} = \blacktriangleleft$, so we get the following




Hence, we see that



Since H has enough points, we observe that the Frobenius condition is satisfied. □

Lemma 3.14. *Let H be a pre-HF algebra; (H, \bullet, \circ) is an integral Hopf algebra if and only if $\square = \curvearrowright$.*


Proof. Note first that $\square = \curvearrowright$ iff $\square^{-1} = \curvearrowleft$. We will use this second form in the following proof. The implication in one direction follows from Lemma 3.6. For the other direction, suppose that $\square^{-1} = \curvearrowleft$. Note that



which implies simply that



Hence, we get




The proof that \circ is a right integral is similar. □

Lemma 3.15. *Let H be a pre-HF algebra such that (H, \bullet, \circ) is an integral Hopf algebra, and let $(\cdot)^\circ$ be the duality defined by the green Frobenius algebra (cf. Lemma 2.5). Then:*



Proof. The first statement is clear from the definition of the green dual and \circ . For the second statement, we see that



where $(*)$ comes from Lemma 3.14. □

Corollary 3.16. *Let H be a pre-HF algebra such that (H, \bullet, \circ) is an integral Hopf algebra. $(H, \blacktriangleright, \blacktriangleleft, \blacktriangleright^\circ, \blacktriangleleft^\circ, \blacktriangle)$ forms a Hopf algebra, where $\blacktriangle = \text{[diagram]}$.*

Proof. First off, let $(\cdot)^\circ$ be the duality defined by the green Frobenius algebra as in Lemma 2.5. Then

$$\blacktriangle^\circ = \text{[diagram]} = \text{[diagram]}$$

It is clear then, that $\blacktriangle = (\blacktriangle^\circ)^{-1}$.

Recall that, by Proposition 2.10, the antipode of H^σ is the inverse of H . We also see that, in a similar manner to Definition 2.11, $(\cdot)^\circ$ will give us another Hopf Algebra, $H^\circ : (H, \blacktriangleright^\circ, \blacktriangleleft^\circ, \blacktriangleright^\circ, \blacktriangleleft^\circ, \blacktriangle^\circ)$. Hence, as H° is a Hopf algebra with antipode \blacktriangle° , and by Lemma 3.15, $(H, \blacktriangleright, \blacktriangleleft, \blacktriangleright^\circ, \blacktriangleleft^\circ, \blacktriangle) = (H^\circ)^\sigma$, so we have our result. \square

Theorem 3.17. *Let H be a Hopf algebra such that the object H has some weak right dual H^* . Then H is a Hopf-Frobenius algebra if and only if H fulfills the Frobenius condition.*

Proof. Given a Hopf algebra that fulfills the Frobenius condition, we see that from Theorem 3.11 and Corollary 3.16 that we get a Hopf-Frobenius algebra.

Suppose that H is a Hopf-Frobenius algebra. Then by Lemma 3.14 the green Hopf algebra is an integral Hopf algebra. From Definition 3.9, we get the following:

$$\text{[diagram]} = \text{[diagram]} = \text{[diagram]} = \text{[diagram]} = \text{[diagram]}$$

\square


Theorem 3.18. *Let H be a pre-HF algebra such that, for some \blacktriangle , $(H, \blacktriangleright, \blacktriangleleft, \blacktriangleright^\circ, \blacktriangleleft^\circ, \blacktriangle)$ forms a Hopf algebra; then H extends to a Hopf-Frobenius algebra.*

Proof. We begin by showing that [diagram] is a right integral and is coinvertible. The proof that


[diagram] is a left cointegral and invertible is similar. First off, by the Frobenius algebra structure of [diagram]


$$\text{[diagram]} = \text{[diagram]} = \text{[diagram]}$$


We see that this map gets copied via [diagram]




This implies that it is coinvertible as follows




We may use similar proofs to show that  is a left cointegral, is group-like and is invertible. To get an integral Hopf algebra from this, observe that



Thus, we may define the following point and copoint, α and β




which have inverses




This means that $((H, \text{node}), \alpha, \beta)$ is an integral Hopf algebra. Since α and β are invertible, Proposition C.8 tells us that we may construct new Frobenius algebras with α and β as unit counit of their respective Frobenius algebras. Let us denote these Frobenius algebras as $(H, \text{node}, \text{node}, \text{node}^\beta, \text{node}^\beta)$ and $(H, \text{node}, \text{node}, \text{node}^\alpha, \text{node}^\alpha)$. We now have a pre-HF algebra, on H such that $(H, \text{node}^\beta, \text{node}^\alpha)$ is an integral Hopf algebra. Hence, by Corollary 3.16, we have a Hopf Frobenius algebra. \square

Lemma 5.2. *The morphism \square is a Hopf algebra homomorphism between H_{\bullet}^{op} and H_{\circ}^**

Proof. We will only show that \square is a homomorphism for node^* , the rest of the structure maps will have similar proofs. We first note that



Hence, we see that



□

B Taft Hopf algebra for $n = 2$

Here we shall state the Hopf-Frobenius algebra for the 4 dimensional Taft Hopf algebra explicitly. It is generated by g and x , and has the structure

| | | | | | | | | | | | |
|------|------|------|-------|------|------|-------------------------------|------|---|------|------|------|
| | 1 | x | g | gx | | 1 | 1 | | 1 | | 1 |
| 1 | 1 | x | g | gx | 1 | $1 \otimes 1$ | 1 | 1 | 1 | 1 | 1 |
| x | x | 0 | $-gx$ | 0 | x | $1 \otimes x + x \otimes g$ | x | 0 | x | gx | gx |
| g | g | gx | 1 | x | g | $g \otimes g$ | g | 1 | g | g | g |
| gx | gx | 0 | $-x$ | 0 | gx | $g \otimes gx + gx \otimes 1$ | gx | 0 | gx | $-x$ | $-x$ |

\mathbf{FVect}_k is a compact closed category, so the integral projection is the map

| | |
|------|---------------|
| | \mathcal{I} |
| 1 | 0 |
| x | $x - gx$ |
| g | 0 |
| gx | 0 |

Hence, the element $x - gx$ is a left cointegral, and the copoint defined as

$$a \mapsto \begin{cases} 1 & \text{if } a = x \\ 0 & \text{otherwise} \end{cases}$$

is a right integral. Hence, by Theorem 3.11, these shall be our unit and counit respectively. It is now possible to construct the resulting Hopf-Frobenius algebra, but we shall explicitly state the structure maps. The green Frobenius algebra is

| | | | | | | | | | | | |
|------|------|------|---|-----|------|------|------|------|------|---|---|
| | $:=$ | | 1 | x | g | gx | | $:=$ | | 1 | $1 \otimes x + gx \otimes g - g \otimes gx + x \otimes 1$ |
| 1 | | 1 | 0 | 1 | 0 | 0 | 1 | | 1 | $1 \otimes x + gx \otimes g - g \otimes gx + x \otimes 1$ | |
| x | | x | 1 | 0 | 0 | 0 | x | | x | $x \otimes x - gx \otimes gx$ | |
| g | | g | 0 | 0 | 0 | 1 | g | | g | $g \otimes x + x \otimes g - 1 \otimes gx + gx \otimes 1$ | |
| gx | | gx | 0 | 0 | -1 | 0 | gx | | gx | $gx \otimes x + x \otimes gx$ | |

| | | | |
|--|------|--|---|
| | $:=$ | | $= 1 \otimes x + gx \otimes g - g \otimes gx + x \otimes 1$ |
|--|------|--|---|

and the red Frobenius algebra

| | | |
|--|----|--|
| | := | |
|--|----|--|

| | | | | | |
|----|---|---|----|----|----|
| | | 1 | x | g | gx |
| 1 | 0 | 0 | 0 | -1 | |
| x | 1 | 0 | 0 | 0 | |
| g | 0 | 1 | 0 | 0 | |
| gx | 0 | 0 | -1 | 0 | |

| | | |
|--|----|--|
| | := | |
|--|----|--|

| | | | | | |
|----|---|---|----|-----|----|
| | | 1 | x | g | gx |
| 1 | 0 | 0 | 0 | -1 | |
| x | 1 | x | 0 | 0 | |
| g | 0 | g | 0 | 0 | |
| gx | 0 | 0 | -g | -gx | |

:= = $1 \otimes x + x \otimes g - g \otimes gx - gx \otimes 1$

C Additional Background Material

In this section we provide additional definitions and basic properties to flesh out the background material of Section 2.

C.1 Categories with duals

Proposition C.1. *In a monoidal category \mathcal{C} suppose that A has two right duals (B_1, d_1, e_1) and (B_2, d_2, e_2) ; then there exists an isomorphism $f : B_1 \cong B_2$, satisfying the equations shown below.*

| | | |
|--|----|--|
| | := | |
|--|----|--|

| | | |
|--|---|--|
| | = | |
|--|---|--|

| | | |
|--|---|--|
| | = | |
|--|---|--|

Proof. Define f as shown above; the required equations follow immediately. □

C.2 Monoids and Comonoids

Definition C.2. A *monoid* in a monoidal category \mathcal{C} consists of an object M , a binary multiplication $\mu : M \otimes M \rightarrow M$ and a unit morphism $\eta : I \rightarrow M$ obeying the familiar associativity and unit laws, shown in diagram form below.

| | | |
|--|---|--|
| | = | |
|--|---|--|

| | | |
|--|---|--|
| | = | |
|--|---|--|

A *comonoid* in \mathcal{C} is a monoid in \mathcal{C}^{op} , concretely depicted below.

| | | |
|--|---|--|
| | = | |
|--|---|--|

| | | |
|--|---|--|
| | = | |
|--|---|--|

A (co)monoid is called (co)commutative if its (co)multiplication is invariant under the exchange map, as depicted below.

| | | |
|--|---|--|
| | = | |
|--|---|--|

| | | |
|--|---|--|
| | = | |
|--|---|--|

In this paper we will *not* assume commutativity or cocommutativity.

Definition C.3. Given a monoid $(M, \begin{smallmatrix} \circlearrowleft \\ \circlearrowright \end{smallmatrix}, \circlearrowright)$, a point $a : I \rightarrow M$ is *left invertible* if there exists a point $l : I \rightarrow M$ satisfying the left equation below; it is *right invertible* if there exists $r : I \rightarrow M$ satisfying the right equation; it is *invertible* if it is both left and right invertible, in which case the two inverses coincide.

$$\begin{array}{c} \triangleleft \quad \triangleleft \\ \downarrow \quad \downarrow \\ l \quad a \\ \downarrow \\ \circlearrowright \end{array} = \begin{array}{c} \circlearrowright \\ \downarrow \end{array} = \begin{array}{c} \triangleleft \quad \triangleleft \\ \downarrow \quad \downarrow \\ a \quad r \\ \downarrow \\ \circlearrowright \end{array}$$

Co-invertibility of co-points $\alpha : M \rightarrow I$ with respect to a comonoid is defined dually.

C.3 Frobenius algebras

In the following lemmas we will assume that we have a given Frobenius algebra $(F, \begin{smallmatrix} \circlearrowleft \\ \circlearrowright \end{smallmatrix}, \circlearrowright, \circlearrowleft, \circlearrowright)$.

Definition C.4. A Frobenius algebra is called *symmetric* if its cap (or equivalently its cup) is invariant under the symmetry.

$$\begin{array}{c} \circlearrowright \\ \downarrow \end{array} = \begin{array}{c} \circlearrowleft \\ \downarrow \end{array} \qquad \begin{array}{c} \circlearrowleft \\ \downarrow \end{array} = \begin{array}{c} \circlearrowright \\ \downarrow \end{array}$$

Lemma C.5. *There is a bijective correspondence between invertible points for the monoid and coinvertible copoints for the comonoid.*

Proof. Let $(\cdot)^\circ$ be the duality induced by the cup and cap; then $u : I \rightarrow F$ is invertible iff and only if $u^\circ : F \rightarrow I$ is coinvertible. \square

Lemma C.6. *Let u be a coinvertible element of the comonoid. Define*

$$\beta(u) := \begin{array}{c} \circlearrowright \\ \downarrow \\ \square \\ u \end{array} \qquad \bar{\beta}(u) := \begin{array}{c} \circlearrowleft \\ \downarrow \\ \square \\ u^{-1} \end{array}$$

Then $\beta(u)$ is a Frobenius form for the monoid $(F, \begin{smallmatrix} \circlearrowleft \\ \circlearrowright \end{smallmatrix}, \circlearrowright)$.

Proof. We must show that the equations of 2.6 hold. The first follows from associativity of the monoid. For the second we have:


$$\begin{array}{c} \circlearrowright \\ \downarrow \\ \square \\ u \end{array} \begin{array}{c} \circlearrowright \\ \downarrow \\ \square \\ u^{-1} \end{array} = \begin{array}{c} \circlearrowright \\ \downarrow \\ \square \\ u \end{array} \begin{array}{c} \circlearrowright \\ \downarrow \\ \square \\ u^{-1} \end{array} = \begin{array}{c} \circlearrowright \\ \downarrow \\ \square \\ u \end{array} \begin{array}{c} \circlearrowright \\ \downarrow \\ \square \\ u^{-1} \end{array} = \begin{array}{c} \circlearrowright \\ \downarrow \\ \square \\ u \end{array} \begin{array}{c} \circlearrowright \\ \downarrow \\ \square \\ u^{-1} \end{array} = \begin{array}{c} \circlearrowright \\ \downarrow \\ \square \\ u \end{array} \begin{array}{c} \circlearrowright \\ \downarrow \\ \square \\ u^{-1} \end{array}$$

and similarly for the other side. Note that $\beta(u) = \beta(v)$ implies $u = v$ by the uniqueness of inverses. \square

Lemma C.7. Suppose that β is a Frobenius form on \mathcal{V} ; then we obtain a coinvertible element $u : F \rightarrow I$ as follows:



Proof. We need only to show that u^{-1} is the coinverse of u .



□

Combining the three preceding lemmas we obtain:

Proposition C.8. *There is a bijective correspondence between the invertible elements of a monoid and the Frobenius forms definable on it.*